

# **SOUTHERN ARIZONA-SOUTHWESTERN NEW MEXICO PROVINCE (025)**

**By W.C. Butler**

## **INTRODUCTION**

This geographically extensive, but unproductive province, covers about 60,900 sq mi from northwestern Arizona to southern Arizona to southwestern New Mexico, primarily in the Basin and Range Physiographic Province, but includes some of the northwest-trending Mogollon transition zone between the Colorado Plateau and the block-faulted terrain. Outcrops in the transition zone are mostly Early and Middle Proterozoic rocks and eroded remnants of Paleozoic and Mesozoic rocks covered by Cenozoic volcanics. Outcrops in the Basin and Range are eroded remnants of similar rocks being buried in their own debris. Except for the southwesternmost part of Arizona, which is geologically similar to the Salton Trough Province 016, the western half of province 025 has very little petroleum potential. Maximum thickness of Phanerozoic rocks is 25,000–30,000 ft in southwesternmost New Mexico.

Province 025 has an extremely complex geologic history that includes several major episodes of Phanerozoic tectonism. Deformation accompanied these province events: (1) Late Paleozoic interplate compression rejuvenated basement faults and formed basins and uplifts on the cratonic platform-shelf; (2) subduction to the southwest caused early- to mid-Mesozoic magmatic arc emplacement and formation of the Mogollon highlands; (3) Early Cretaceous rifting (Bisbee Basin, that is, the northern extension of the Chihuahua trough-aulacogen); (4) Late Mesozoic to Eocene compression produced basement-cored uplifts and localized northeast-directed thrusting; and, (5) middle to late Cenozoic extension caused block and detachment faulting.

Precambrian, Paleozoic, Mesozoic, and Tertiary rocks are exposed in fault-block mountains surrounded by Neogene and Quaternary basin-fill alluvium. Metamorphic core complexes, plutons and other igneous intrusions, and world-class Laramide ore deposits occur within the area. For additional information see Butler (1989) and Jenny and Reynolds (1989a).

Evidence suggests that during the Precambrian a major (100-mi-wide), but enigmatic and controversial, west-northwest-trending shear zone, the Texas lineament (Muehlberger, 1980), may have existed between what is now the site of El Paso, Texas, and the present Pacific Ocean. It may be the longest zone of weakness, or wrench fault

(Sylvester, 1984), in the northern hemisphere. Movement in opposite senses probably occurred intermittently from Precambrian through Laramide times. Transpressional uplifts or anticlinal welts, tilt blocks, drag folds, local thrusts and normal faults, and pull-apart basins along slight changes in direction of the fault are probably present but obscured by later tectonism. Hydrocarbon traps may have been formed as a result of these structures. Examples of crustal sags associated with the shear zone might include the Late Paleozoic marine Pedregosa Basin and the area of southwestern Arizona where as much as 25,000 ft of the nonmarine Jurassic or Early Cretaceous McCoy Mountains Formation (Harding and Coney, 1985) was deposited. Other transform-like strike-slip movement has been documented on the Jurassic Mojave-Sonoran megashear (northern Mexico); it may be en echelon with the Texas lineament.

Paleozoic shelf sediments were deposited on a passive plate margin near the southwestern terminus of the transcontinental arch. Late Paleozoic petroleum source and reservoir strata were deposited in the sometimes-restricted Pedregosa Basin of southeastern Arizona and southwestern New Mexico. Figure 2 illustrates the stratigraphic column for the Pedregosa Basin. For a very brief tectonic and depositional history, see the discussion under Northern Arizona, Province 024.

In the eastern half of this province, the Triassic and Jurassic Periods were generally times of erosion and intrusive and extrusive igneous activity. Also, as much as 15,000 ft of Early and Late Cretaceous sediments accumulated in the Bisbee Basin; most were poorly sorted clastics, but Early Cretaceous basin-margin carbonate algal-rudist reefs did develop. Generalized Late Cretaceous depositional settings included alluvial fans adjacent to a coastal plain and a fluvial-marine deltaic complex.

Four speculative plays are considered in this province: two in the Pennsylvanian-Permian section of the Pedregosa Basin, one play in seven of the nonmarine Tertiary basins of southeastern Arizona, and one marine play of Neogene age in the Altar Basin (proto-Gulf of California) of southwesternmost Arizona. They are the Alamo-Hueco Basin Play (2501), Pedregosa Basin Play (2502), Seven Tertiary continental Basins Play (2503), and Altar-San Luis Play (2504). Exploration intensity has been light with approximately one borehole drilled per 325 sq mi.

## **ACKNOWLEDGMENTS**

Scientists affiliated with the American Association of Petroleum Geologists and from various State geological surveys contributed significantly to play concepts and definitions. Their contributions are gratefully acknowledged.

## CONVENTIONAL PLAYS

### 2501. ALAMO HUECO BASIN PLAY (HYPOTHETICAL)

This hypothetical oil and nonassociated gas play is defined on Pennsylvanian and Early Permian source and reservoir rocks in the deeper part of the Pedregosa Basin of southwestern New Mexico. Evidence comes from the key stratigraphic section of the Big Hatchet Mountains (Wengerd, 1970; Thompson and Jacka, 1981) near the center of the play.

**Reservoirs:** Reservoirs might include fractured shales of the Horquilla Limestone. Sandy fluvial to deltaic clastic units (moderately sorted quartzarenite) of the 1,000-ft-thick Earp Formation have reservoir characteristics and may be sealed by Earp Formation micrites, reddish shales, and calcite-cemented siltstones, or by the superjacent Early Permian shales, impermeable carbonates, or gypsum beds. The Earp Formation is probably lithologically correlative with the gas-productive Abo Formation in southeastern New Mexico.

**Source rocks:** Potential source rocks are black, 2,500–3,500-ft thick shales, mudstones, dark limestones, and reef carbonates of the Horquilla Limestone. The deep marine basinal mudstones probably reached thermal maturity during the Cretaceous Period and generated oil that migrated northward updip into coeval shallow-marine shelf dolomite facies (porous phylloid algal mounds). Vitrinite reflectance data from Paleozoic outcrops in southwestern New Mexico range from about  $R_o$  0.6 to 1.8 percent. Thermal maturity increases southeasterly through the play into northern Mexico toward deeper pre-Laramide burial sites.

**Traps:** In addition to the stratigraphic traps within the Horquilla, as noted above, the various styles and times of complex deformation provide structural traps. An outline of tectonic events and associated structures is noted in the introductory section. Half-grabens, basement-cored uplifts, northeast-directed Laramide overthrusts, reverse and normal faults, right-lateral strike-slip wrench faults, and major northwest-trending folds are some of the potential trapping mechanisms in this play. The Paleozoic strata, which are 3,000 – 14,000 ft thick north to south, respectively, may have several thrust-repeated sections in the southern part. In the northern part, both the Paleozoic and Cretaceous sections may be involved in numerous thrust sheets. Drilling depths may be as much as 25,000 ft. Although gas is a stable hydrocarbon species in deep parts of this play, it may not have accumulated; that is, the gas-prone areas may have leaked due to the highly

fractured and faulted reservoirs. Tectonic overprinting may not only have created traps, but also may have destroyed them, leaving few large accumulations possible.

**Exploration status and resource potential:** Few of the suspected deep traps have been drilled. Density of exploratory drilling is about one well per 40 sq mi; in the entire Pedregosa Basin this density decreases to one well per 1,340 sq mi for wells that have encountered either the Paleozoic or Precambrian rocks. Twelve boreholes in this play had significant oil and gas shows, and one, Humble Oil No. 1 State BA (TD 14,585 ft) had nearly commercial amounts of gas from the Permian Epitaph Dolomite.

Limiting factors to upgrading the petroleum potential include possible flushing of reservoirs by meteoric water and the rupturing of traps and seals by repeated post-migration Neogene Basin-and-Range faulting. Thermal destruction of hydrocarbons may have occurred in this highly-mineralized terrain close to plutons and volcanic cauldrons; however, the thermal effects of intrusions in some cases may actually mature the source rock and are not thought to severely alter the host rock more than a few tens of feet away from its contact with hot igneous rock.

## **2502. PEDREGOSA BASIN PLAY (HYPOTHETICAL)**

This hypothetical, highly-speculative, oil and nonassociated gas play is in the Basin and Range Physiographic Province of southeastern Arizona. The play boundary is defined by thick upper Paleozoic strata (4,000–4,750 ft) that may contain some source rocks within the thermal zone of oil generation in the proximal part of the Pedregosa Basin. Outcrops in the horsts surrounding the play are over-mature with respect to oil.

**Reservoirs:** Potential reservoir rocks primarily include the Horquilla Limestone carbonates (see play 2501) and minor sandstones totaling on average of 1,000 ft thick for the formation, plus other intercalated upper Paleozoic marine shelf carbonates and tidal flat and deltaic siltstones and sandstones, for example, the Earp Formation totaling on average of 1,200 ft thick. The carbonates locally have vuggy porosity. Reservoirs may be sealed by upper Paleozoic units, such as the massive gypsum beds of the Epitaph Dolomite, or dense carbonates or fine-grained intercalated clastics below the Scherrer Formation, or the orthoquartzites of the Scherrer. Diagenetic porosity pinchouts between the various carbonate facies of the Colina and Epitaph may also prevent migration.

**Source rocks:** Potential source rocks of fair to average quality could include the dark carbonates of the Mississippian Escabrosa Limestone and Paradise Formation, the

Pennsylvanian carbonates and thin shales of the Black Prince Limestone and Horquilla Limestone, and the dark Permian shelf limestones, dolomites, and shales in the Earp Formation, Colina Limestone, Epitaph Dolomite, and Concha and Rainvalley Formations. The Pennsylvanian-Permian fossiliferous carbonates are consistently dark on fresh surfaces; some of these are biostromal and stromatolitic and produce thick oily residues upon dissolution in acid. Shows of oil and gas have been detected in all of the rock systems in the Pedregosa Basin; the ratio of shows to dry holes is greater than 1.0. Fetid odors are very common in outcrops of this play.

**Timing and migration:** Timing of generation and migration of hydrocarbons are unknown but may have been during early Cretaceous or Laramide time.

**Traps:** Various stratigraphic and structural traps are commonplace; the general types are listed in the province introduction. Unconformities as depicted in the stratigraphic section present an additional trap and seal potential. Targets are expected at 6,000- to 17,000-ft drilling depths due the extreme structural complexity of the area. This complexity may also be responsible for trap destruction.

**Exploration status and resource potential:** Drilling density in this play is about one petroleum-test borehole per 150 sq mi, and only one borehole is deeper than 10,000 ft.; there are only a few tests between 5,000 and 10,000 ft. The potential of this frontier area has been inadequately tested.

### **2503. SEVEN TERTIARY CONTINENTAL BASINS PLAY (HYPOTHETICAL)**

This hypothetical, mostly nonassociated gas play is in the Basin and Range Physiographic Province of southeastern Arizona. The seven basins included are Tucson (Santa Cruz Valley), Sonoita-Cienega, San Simon (Safford Valley), Sulphur Springs (north), Sulphur Springs south (Elfrida), Teran (San Pedro north), and San Pedro (south).

Crustal relaxation and extension, which began in the Oligocene and was most pronounced during the mid-Tertiary, created a general horst and graben (listric block-faulted) terrane. Intermittent uplifting eroded the horsts, and the internally drained basins filled with thick sequences of continental alluvial-fluvial conglomerates, various pyroclastic and flow volcanics, and lacustrine sediments. Rapidly rising horsts shed thousands of feet of very coarse (boulder conglomerates) to very fine clastics, (for example, mudstones) into the asymmetrical grabens and half-grabens.

The lacustrine deposits include evaporites of gypsum, anhydrite, and halite. These mostly Miocene and Pliocene salt masses have been estimated to be from 3,600 ft to more than 12,000 ft thick in the major Tertiary basins of the Arizona Basin and Range Physiographic Province. In the basins of south-central Arizona near Phoenix, they may be as much as 10,000 ft thick. Salt thickness is purely conjectural for southeastern Arizona; the tops of salt masses here may be from 500 to 3,000 ft below the surface. Depths to bedrock are about 4,000 ft to more than 12,000 ft (Oppenheimer and Sumner, 1980). Elevations of the Precambrian basement, that is, the graben floors, are about 6,000 – 10,000 ft below sea level.

The simple model of petroliferous lacustrine deposits, including stomatolitic carbonates, marly limestones, and various chemical precipitates, being rapidly covered by the clastic (reservoir) units and impermeable evaporites and clays (seals) as the lake shorelines shifted, serves as the basis for this play. Seven Tertiary basins, some having connections at the shallower depths, are included in this play; more basins in the Arizona Basin and Range Physiographic Province, may also have petroleum potential, but additional study will be necessary in order to include them.

**Reservoirs, source rocks, and traps:** Source and reservoir rocks are closely associated. Most of the Tertiary Pantano Group is clastic basin-fill; however, it is highly variable and may contain lacustrine deposits (organic-rich clays, evaporites, and carbonates) that could qualify as source rocks. Hypersalinity in the basinal lakes may have preserved organic matter that is possibly of types I, II, and III. These basins exhibit normal to anomalously high heat flow, plutonic and volcanic rocks, and thermal springs; although nonassociated gas is expected, including some biogenic gas, some oil may be present if maturation was augmented by igneous activity.

Environments of deposition in this play during the mid-Tertiary include pediment surfaces (for example, gravel) and alluvial fans near the play edges, and floodplain, fluvial, paludal, terrace, playa, subaerial dune, lake delta, and evaporating pan environments more distal to the source areas. These depositional facies, as well as the abundant layered volcanic units, change rapidly over short distances and could provide not only good reservoirs but also the porosity and permeability pinchouts necessary for trapping. Salt beds intertonguing with the terrestrial clastics may have also trapped migrating petroleum. Unconformable stratigraphic contacts are common and may only very locally prevent migration.

The role of diapirism in the salt masses is essentially unknown and currently indeterminable as to whether it may have caused structural traps to form in deformed beds. Targets are a few hundred feet deep to possibly more than 12,500 ft deep along basin axes. All potential petroleum reservoirs in the seven basins have a high probability of being flushed with meteoric water and fresh water is being heavily mined by relatively shallow water wells in some areas to support the needs of a rapidly increasing population and the agricultural and industrial economy.

**Exploration status and resource potential:** Several dozens of minor hydrocarbon shows are documented in the driller's logs of these basins; most are reported in water wells. More than a dozen minor oil seeps have been reported. Drilling density in this play is about one borehole per 70 sq mi, but only one petroleum borehole per every 410 sq mi. Only four boreholes have been drilled deeper than 5,000 ft in this play. This play depends on exceptionally unique geologic conditions to make it work, and therefore it has only fair hydrocarbon potential.

#### **2504. ALTAR-SAN LUIS BASIN PLAY (HYPOTHETICAL)**

This hypothetical oil and nonassociated gas play is in the Basin and Range Physiographic Province of southwesternmost Arizona (Yuma Desert). The play is defined by very significant onshore and offshore petroleum shows and possible discoveries in Cenozoic sand-shale sequences just to the south in northwestern Sonora, Mexico.

**Source rocks:** Transgressive Cenozoic (mostly Miocene and Pliocene) marine to brackish-water estuarine sediments were deposited in the deep, narrow, euxinic proto-Gulf of California along the present-day axis of the Colorado River (Armentrout and others, 1979). The shales and carbonates, for example, the Pliocene Bouse Formation, may contain as much as 10 percent total organic carbon in places (Guzman, 1981). Highly variable thermal maturities and hydrocarbon types suggest that gas, gas condensate, and oil can all be expected to occur; there may be some biogenic gas in the shallower parts of the play. The Exxon #1 Yuma-Federal well was drilled in 1973 to a depth of 11,444 ft; vitrinite reflectance measurement at TD was  $R_0$  1.1 percent. Known geothermal resource areas are present just a few miles west of this play in the Imperial Valley-Salton Sea area. This close proximity may expedite the maturation of the shallower source rocks nearest the California border.

**Reservoirs:** Reservoir rocks of the Bouse Formation (about 1,000–2,000 ft thick) are within an admixed package of bedded volcanics, carbonates, and turbidites, (conglomerates, sandstones, siltstones, shales); the several thousand feet of Miocene clastics is the secondary target. Miocene evaporites and Neogene fine-grained clastics may provide reservoir seals.

**Traps:** Guzman (1981) suggested that the Altar Basin is genetically related to the oil-rich Los Angeles Basin and other Miocene basins of southern California, but offset by right-lateral movement along the San Andreas transform fault. Drag folding and push-up structures along this strike-slip fault system may have created hydrocarbon traps (see introductory section regarding Texas lineament structures). Facies changes over short distances, and hence porosity-permeability pinchouts are to be expected in this trough, which received a rapid accumulation of sediments. Fault traps may be present. Lastly, mid-Tertiary tilting creating angular unconformities, and erosion creating interformational disconformities, may also enhance petroleum trapping and accumulation. Generation and migration of hydrocarbons is probably still occurring.

#### **Exploration status:**

Seismic refraction data in the northern Gulf of California indicate that unconsolidated Pliocene(?) and Pleistocene sediments may be 5,000 ft thick; semi-consolidated Miocene(?) and Paleogene sediments may be 6,500 ft thick. Gravity studies indicate that Neogene sediments may be 10,000–15,000 ft thick between the Salton Sea and the northern Gulf of California. Depth to Cretaceous bedrock is probably 11,000–13,000 ft in the play.

Miocene gas, possibly commercial, was discovered at a depth of about 13,500 ft as reported in 1981 (see Nations and others, 1983a, for references to the original information) in the northern Gulf of California area 14 mi offshore. Either another discovery may have been made in 1982, or else confirmation and development wells may have been drilled near the 1981 discovery. Accurate information is difficult to obtain, but existing information indicates that the 1981 (or 1982 wells) encountered deltaic sediments of the Colorado River and yielded a gas and condensate discovery in a Pliocene-Pleistocene section 13,000–19,700 ft thick, gas and oil flowed at almost 6 MMCFGPD and 30 BOPD. Other Pemex discoveries have been reported in 1983, 8 mi south of this play. In 1985 gas zones and salt beds were drilled in 16,400 ft of Neogene strata about 30 mi south of the southeast tip of this play in Sonora, Mexico; water production caused abandonment of the well. About two dozen petroleum boreholes,



some having shows, have been drilled in the Yuma area of the United States; five of these are more than 5,000 ft deep.

## **UNCONVENTIONAL PLAYS**

There are no unconventional plays described in this province report. However, unconventional plays listed in the surrounding provinces may include parts of this province. Individual unconventional plays are usually discussed under the province in which the play is principally located.

## REFERENCES

- Aiken, C.L.V., Laughlin, A.W., and West, F.G., 1978, Residual Bouguer gravity anomaly map of New Mexico: University of California Los Alamos Scientific Laboratory LA-7466-map/UC-11, scale 1:500,000.
- Aiken, C.L.V., and Sumner, J.S., 1974, A geophysical and geological investigation of potentially favorable areas for petroleum exploration in southeastern Arizona: Arizona Oil and Gas Conservation Commission Report of Investigations 3, 40 p.
- Allin, D.L., 1990, Colorado Plateau subsurface water flow key: Oil and Gas Journal, v. 88, no. 30, July 23, p. 52-54.
- American Geophysical Union, 1989, Sedimentation and tectonics of western North America, v. 3: Washington, D.C., American Geophysical Union, 28th International Geological Congress, 477 p.
- Anderson, O.J., Lucas, S.G., Love, D.W., and Cather, S.M., eds., 1989, Southeastern Colorado Plateau: New Mexico Geological Society, 40th Field Conference Guidebook, 348 p.
- Anderson, R.Y., and Harshbarger, J.W., eds., 1958, Black Mesa Basin--Guide book of northeastern Arizona: Arizona Geological Society, 9th Field Conference, 205 p.
- Anschutz, P.F., 1980, The overthrust belt--Will it double U.S. gas reserves?: World Oil, v. 190, no. 1, p. 111-116.
- Arizona Oil and Gas Conservation Commission, 1961, Oil, gas and helium in Arizona: Phoenix, Arizona, Arizona Development Board, 108 p.
- Arizona Oil and Gas Conservation Commission, 1987, State of Arizona well location map four: Phoenix, Arizona, August, 38 p.
- Arizona Oil and Gas Conservation Commission, 1992, Arizona well location map and report: Arizona Geological Survey Oil and Gas Publication OG-12, 22 p., scale 1:500,000.
- Armentrout, J.M., Cole, M.R., and TerBest, Harry, Jr., eds., 1979, Cenozoic paleogeography of the western United States: Society of Economic Paleontologists and Mineralogists, Pacific Coast Section Symposium 3, 335 p.
- Armstrong, A.K., Kottowski, F.E., Stewart, W.J., Mamet, B.L., Baltz, E.H., Siemers, W.T., and Thompson, Sam, III, 1979, The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States--New Mexico: U.S. Geological Survey Professional Paper 1110-W, 27 p.

- Arnold, E.C., and Hill, J.M., comps., 1981, New Mexico's energy resources '80: New Mexico Bureau of Mines and Mineral Resources Circular 181, 59 p.
- Baars, D.L., ed., 1979, Permianland: Four Corners Geological Society Guidebook, 9th Field Conference and Field Symposium, 186 p.
- Baars, D.L., 1983, The Colorado Plateau--A geologic history: Albuquerque, New Mexico, University of New Mexico Press, 279 p.
- Bahr, C.W., 1962, The Holbrook anticline, Navajo County, Arizona, *in* Weber, R.H., and Peirce, H.W., eds., Guidebook of the Mogollon Rim region east-central Arizona: New Mexico Geological Society, 13th Field Conference, p. 118-122.
- Barwin, J.R., 1969, Oil possibilities in Arizona: Oil and Gas Journal, v. 67, no. 35, p. 186-188.
- Barwin, J.R., King, R.W., and Hassenfratz, C.A., 1971, Future oil and gas potential of northeast Arizona, *in* Cram, I.H., ed., Future petroleum provinces of the United States--Their geology and potential: American Association of Petroleum Geologists Memoir 15, p. 449-469.
- Bayer, K.C., comp., 1983, Generalized structural, lithologic and physiographic provinces in the fold and thrust belts of the United States: U.S. Geological Survey map, scale 1:2,500,000.
- Bayley, R.W., and Muehlberger, W.R., 1968, Basement rock map of the United States: U.S. Geological Survey Map, scale 1:2,500,000.
- Bayliss, G.S., and Schwarzer, R.R., 1992, New Mexico hydrocarbon source rock evaluation project--New Well Peak section in Big Hatchet Mountains, Hidalgo County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Open-File Report 379, (revised January, 1992), unpaginated.
- Beal, C.H., 1948, reconnaissance of the geology and oil possibilities of Baja California, Mexico: Geological Society of America Memoir 31, 138 p.
- Beatty, Barbara, and Wilkinson, P.A.K., eds., 1986, Frontiers in geology and ore deposits of Arizona and the Southwest: Arizona Geological Society Digest, V. XVI, 554 p.
- Beikman, H.M., Peterson, J.A., Huber, D.F., and Butler, W.C., 1986, Metallic mineral and mineral-fuel resource potential map of Arizona showing major mineral deposits: U.S. Geological Survey Map Mineral Investigations Resource Map MR-94, scale 1:1,000,000.

- Beus, S.S., 1989, Devonian and Mississippian geology of Arizona, *in* Jenney, J.P., and Reynolds, S.J., eds., *Geologic evolution of Arizona: Arizona Geological Society Digest*, V. 17, p. 287-311.
- Billingsley, G.H., 1978, A synopsis of stratigraphy in the western Grand Canyon: Flagstaff, Arizona, Museum of Northern Arizona Research Paper 16, 26 p.
- Bilodeau, W.L., and Lindberg, F.A., 1983, Early Cretaceous tectonics and sedimentation in southern Arizona, southwestern New Mexico and northern Sonora, Mexico, *in* Reynolds, M.W., and Dolly, E.D., eds., *Mesozoic paleogeography of west-central United States: Society of Economic Mineralogists and Paleontologists, Symposium 2*, p. 173-188.
- Blackwell, D.D., 1978, Heat flow and energy loss in the western United States, *in* Smith, R.B., and Eaton, G.P., eds., *Cenozoic tectonics and regional geophysics of the Western Cordillera: Geological Society of America Memoir 152*, p. 175-208.
- Blakey, R.C., 1979, Oil impregnated carbonate rocks of the Timpweap Member Moenkopi Formation, Hurricane Cliffs area, Utah and Arizona: *Utah Geology*, v. 6, p. 45-53.
- Brady, B.T., 1984, Selected geologic and hydrologic characteristics of the Basin and Range Province, western United States, coal, oil and gas wells, seeps and tar sandstone occurrences: U.S. Geological Survey Miscellaneous Investigations Series Map I-1522-E, scale 1:2,500,000.
- Branson, C.C., ed., 1962, Pennsylvanian system in the United States--A symposium: American Association of Petroleum Geologists, 508 p.
- Brennan, D.J., 1989, Hydrocarbon possibilities in Yuma area, Arizona [abs.]: *American Association of Petroleum Geologists Bulletin*, v. 73, no. 3, p. 337.
- Broadhead, R.F., and Black, B.A., 1989, Petroleum exploration wells drilled in west-central New Mexico, *in* Anderson, O.J., and others, eds., *South eastern Colorado Plateau: New Mexico Geological Society, 40th Field Conference Guidebook*, p. 287-296.
- Bryant, D.L., 1968, Diagnostic characteristics of the Paleozoic formations of southeastern Arizona, *in* Titley, S.R., ed., *Southern Arizona guidebook III: Tucson, Arizona, Arizona Geological Society*, p. 33-47.
- Butler, W.C., 1971, Permian sedimentary environments in southeastern Arizona, *in* McCullough, E.J., ed., *Arizona Geological Society Digest*, V. IX, p. 71-94.

- Butler, W.C., 1985, Petroleum potential of the Hualapai Indian Reservation in northwestern Arizona--A preliminary qualitative assessment for the tribe: unpublished U.S. Geological Survey and Bureau of Indian Affairs Administrative Report, on file at Bureau of Indian Affairs, 20 p.
- Butler, W.C., 1988a, The rationale for assessment of undiscovered, economically recoverable oil and gas in central and northern Arizona Play analyses of seven favorable areas: U.S. Geological Survey Open-File Report 87-450-V, 145 p.
- Butler, W.C., 1988b, The rationale for assessment of undiscovered, economically recoverable oil and gas in south-central New Mexico--A geologic overview and play analysis of two favorable areas: U.S. Geological Survey Open-File Report 88-450-B, 134 p.
- Butler, W.C., 1989a, The geologic setting of southern Arizona and southwestern New Mexico, with a rationale for assessment of undiscovered, economically recoverable oil and gas--A summary of four potential plays: U.S. Geological Survey Open-File Report 88-450-M, 150 p.
- Butler, W.C., 1989b, Strategic crude oil storage in Arizona evaporite masses--An initial description, evaluation, and selected references of four potential sites: unpublished U.S. Geological Survey Administrative Report, on file with the Branch of Petroleum Geology, Denver, Colorado, 109 p.
- Butler, W.C., 1992, Elevation data for the Precambrian surface in the central and southern Colorado Plateau and vicinity: U.S. Geological Survey Open File Report 92-194, 69 p.
- Callender, J.F., Wilt, J.C., and Clemons, R.E., eds., 1978, Land of Cochise Southeastern Arizona: New Mexico Geological Society Guidebook, 29th Field Conference, 372 p.
- Cather, S.M., and Johnson, B.D., 1984, Eocene tectonics and depositional setting of west-central New Mexico and eastern Arizona: New Mexico Bureau of Mines and Mineral Resources Circular 192, 33 p.
- Chapin, C.E., ed., 1983, Socorro region II: New Mexico Geological Society, 34th Field Conference Guidebook, 344 p.
- Christiansen, R.L., and Lipman, P.W., 1972, Cenozoic volcanism and plate tectonic evolution of the western United States, II. Late Cenozoic: Philosophical Transactions of the Royal Society of London, Series A, v. 271, no. 1213, p. 249-284.
- Clemons, R.E., King, W.E., and Mack, G.H., eds., 1986, Truth or Consequences region: New Mexico Geological Society, 37th Field Conference Guidebook.

- Clemons, R.E., and Lawton, T.F., 1991, Petrographic and well log analysis of four exploration wells in southwestern New Mexico: New Mexico Bureau of Mines and Mineral Technology Open-File Report 381, 97 p.
- Condie, K.C., 1981, Precambrian rocks of the southwestern United States and adjacent areas of Mexico: New Mexico Bureau of Mines and Mineral Resources, Resource Map 13, scale 1:500,000.
- Condie, K.C., 1982, Plate-tectonics model for Proterozoic continental accretion in the Southwestern United States: *Geology*, v. 10, p. 37-42.
- Coney, P.J., 1979, Tertiary evolution of Cordilleran metamorphic core complexes, *in* Armentrout, J.M., Cole, M.R., and TerBest, H., Jr., eds., *Cenozoic paleogeography of the western United States*: Society of Economic Paleontologists and Mineralogists, Symposium 3, p. 15-28.
- Conley, J.N., and Giardina, S., Jr., 1979, Favorable and potentially favorable areas for hydrocarbon and geothermal energy sources in northeastern Arizona: Arizona Oil and Gas Conservation Commission, Report of Investigations 7, 56 p.
- Corbitt, L.L., and Woodward, L.A., 1973, Tectonic framework of Cordilleran foldbelt in southwestern New Mexico: *American Association of Petroleum Geologists Bulletin*, v. 57, p. 2207-2216.
- Cordoba, D.A., Wengerd, S.A., and Shomaker, John, eds., 1969, *Guidebook of the Border Region*: New Mexico Geological Society, 20th Field Conference, 218 p.
- Crysdale, B.L., and Schenk, C.J., 1988, Bitumen-bearing deposits of the United States: *U.S. Geological Survey Bulletin* 1874, 45 p.
- Dane, C.H., and Bachman, G.O., 1965, Geologic map of New Mexico: U.S. Geological Survey and New Mexico Bureau of Mines and Mineral Resources, scale 1:500,000.
- Davis, G.H., 1978, Monocline fold pattern of the Colorado Plateau, *in* Matthews, V., III, ed., *Laramide folding associated with basement block faulting in the western United States*: Geological Society of America Memoir 151, p. 215-233.
- Davis, G.H., 1979, Laramide folding and faulting in southeastern Arizona: *American Journal of Science*, v. 279, p. 543-569.
- Davis, G.H., and VandenDolder, E.M., eds., 1987, Geologic diversity of Arizona and its margins--Excursions to choice areas: Arizona Bureau of Geology and Mineral Technology, Geological Survey Branch Special Paper 5, 422 p.

- DeFord, R.K., and Kehle, R.O., Chairmen, 1976, Geothermal gradient map of North America: American Association of Petroleum Geologists and U.S. Geological Survey, scale 1:5,000,000.
- Desborough, G.A., Poole, F.G., Daws, T.A., and Scarborough, R.B., 1984, Hydrocarbon source rock evaluation of the Middle Proterozoic Apache Group Gila County Arizona, *in* Woodward, J., and others, eds., Hydrocarbon source rocks of the Greater Rocky Mountain region: Rocky Mountain Association of Geologists Symposium, p. 51-55.
- Dickinson, W.R., 1981, Plate tectonic evolution of the southern Cordillera, *in* Dickinson, W.R., and Payne, W.D., eds., Relations of tectonics to ore deposits in the southern Cordillera: Arizona Geological Society Digest, V. XIV, p. 113-135.
- Dickinson, W.R., 1989, Tectonic setting of Arizona through geologic time, *in* Jenney, J.P., and Reynolds, S.J., eds., Geologic evolution of Arizona: Arizona Geological Society Digest, V. 17, p. 1-16.
- Dickinson, W.R., and Klute, M.A., eds., 1987, Mesozoic rocks of southern Arizona and adjacent areas: Arizona Geological Society Digest, V. 18, 394 p.
- Dixon, G.H., Baltz, D.H., and Stipp, T.F., 1954, Records of wells drilled for oil and gas in New Mexico: U.S. Geological Survey Circular 33, 79 p.
- Drewes, H.D., 1981, Tectonics of southeastern Arizona: U.S. Geological Survey Professional Paper 1144, 96 p.
- Eberly, L.D., and Stanley, T.B., Jr., 1978, Cenozoic stratigraphy and geologic history of southwestern Arizona: Geological Society of America Bulletin, v. 89, p. 921-940.
- Elston, D.P., 1979, Late Precambrian Sixtymile Formation and orogeny at top of the Grand Canyon Supergroup, northern Arizona: U.S. Geological Survey Professional Paper 1092, 20 p.
- Elston, D.P., 1989, Grand Canyon Supergroup, northern Arizona--Stratigraphic summary and preliminary paleomagnetic correlations with parts of other North American Proterozoic successions, *in* Jenney, J.P., and Reynolds, S.J., eds., Geologic evolution of Arizona: Arizona Geological Society Digest, V. 17, p. 259-272.
- Fassett, J.E., ed., 1978, Oil and gas fields of the Four Corners area, v. I and II: Four Corners Geological Society, 727 p.
- Foster, R.W., 1964, Stratigraphy and petroleum possibilities of Catron County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Bulletin 85, 55 p.



- Foster, R.W., 1978, Oil and gas evaluation of White Sands Missile Range and Fort Bliss Military Reservation, south-central New Mexico: New Mexico Bureau of Mines and Mineral Resources, Open-File Report 92, 60 p.
- Fouch, T.D., and Magathan, E.R., eds., 1980, Paleozoic paleogeography of the west-central United States: Society of Economic Paleontologists and Mineralogists, Symposium 1, 431 p.
- Frazier, W.J., and Schwimmer, D.R., 1987, Regional stratigraphy of North America: New York, New York, Plenum Press, 719 p.
- Fritz, M., 1990, Precambrian source sparks imaginations--Chuar inspires quest in the West: American Association of Petroleum Geologists Explorer, October, p. 8-9.
- Frost, E.G., and Martin, D.L., eds., 1982, Mesozoic-Cenozoic tectonic evolution of the Colorado River Region, California, Arizona, and Nevada: San Diego, California, Cordilleran Publishers, Inc., Anderson-Hamilton Volume, (published in conjunction with a symposium and field trip for the Geological Society of America), 608 p.
- Gehrels, G.E., and Spencer, J.E., eds., 1990, Geologic excursions through the Sonoran Desert region, Arizona and Sonora: Arizona Geological Society Special Paper 7, Field-Trip Guidebook, Cordilleran Section, Geological Society of America, 202 p.
- Gerrard, T.A., 1966, Environmental studies of Fort Apache Member, Supai Formation, east-central Arizona: American Association of Petroleum Geologists Bulletin, v. 50, p. 2434-2463.
- Giardina, S., Jr., 1979, Geologic review of northwestern Arizona for petroleum exploration investigators: Arizona Oil and Gas Conservation Commission, Special Publication No. 4, 72 p.
- Grant, P.R., Jr., and Foster, R.W., 1989, Future petroleum provinces in New Mexico--Discovering new reserves: New Mexico Bureau of Mines and Mineral Resources, 94 p. (17" x 22" folio), prepared for the New Mexico Research and Development Institute.
- Greenwood, E., Kottlowski, F.E., and Thompson, S., III, 1977, Petroleum potential and stratigraphy of Pedregosa Basin--Comparison with Permian and Orogrande Basins: American Association of Petroleum Geologists Bulletin, v. 61, p. 1448-1469.
- Grim, P.J., and Berry, G.W., 1979, Geothermal energy in the western United States, *in* Muffler, L.J.P., ed., Assessment of geothermal resources of the United States--1978: National Oceanic and Atmospheric Administration, National Geophysics Solar-Terrestrial Data Center and U.S. Geological Survey Circular 790, scale 1:2,500,000.

- Gurgel, K.D., ed., 1983, Energy resources map of Utah: Utah Geological and Mineral Survey Map 68, scale 1:500,000.
- Gustafson, V.O., 1981, Petroleum geology of the Devonian and Mississippian rocks of the Four Corners region, *in* Wiegand, D.L., ed., Geology of the Paradox Basin: Rocky Mountain Association of Geologists Field Conference Guidebook, p. 101-109.
- Guzman, A.B., 1981, Petroleum prospects in Mexico's Altar Desert: Oil and Gas Journal, v. 79, no. 34, August 24th, p. 110-126.
- Hamblin, W.K., 1965, Origin of "reverse drag" on the downthrown side of normal faults: Geological Society of America Bulletin, v. 76, p. 1145-1164.
- Harding, L.E., and Coney, P.J., 1985, The geology of the McCoy Mountains Formation, southeastern California and southwestern Arizona: Geological Society of America Bulletin, v. 96, p. 755-769.
- Harrison, G.C., 1976, Facies analysis of the Devonian of the Black Mesa Basin, Arizona: Flagstaff, Arizona, Northern Arizona University, unpublished M.S. thesis, 57 p.
- Hawley, J.W., and Lozinsky, R.P., 1992, Hydrogeologic framework of the Mesilla Basin in New Mexico and western Texas: New Mexico Bureau of Mines and Mineral Resources, Open-File Report 323, 55 p.
- Hayes, P.T., 1970, Cretaceous paleogeography of southern Arizona and adjacent areas: U.S. Geological Survey Professional Paper 658-B, 42 p.
- Hayes, P.T., 1975, Cambrian and Ordovician rocks of southern Arizona and New Mexico and westernmost Texas: U.S. Geological Survey Professional Paper 873, 98 p.
- Hereford, Richard, 1977, Deposition of the Tapeats Sandstone (Cambrian) in central Arizona: Geological Society of America Bulletin, v. 88, p. 199-211.
- Heylman, E.B., 1960, Kanab--The oil signs are there: Oil and Gas Journal, v. 58, no. 24, p. 168-171.
- Heylman, E.B., 1981, Geology of promising Holbrook Basin: Oil and Gas Journal, v. 79, p. 117-118, 123.
- Hills, J.M., and Kottowski, F.E., coordinators, 1983, Correlation of stratigraphic units in North America--Southwest/southwest mid-continent correlation chart, Lindberg, F.A., ed.: American Association of Petroleum Geologists COSUNYA chart.

- Hintze, L.F., comp., 1980, Geologic map of Utah: Utah Geologic and Mineralogic Survey, scale 1:500,000.
- Hintze, L.F., 1988, Geologic history of Utah--A field guide to Utah's rocks: Brigham Young University Geology Studies Special Publication 7, 204 p.
- Huddle, J.W., and Dobrovolsky, E., 1952, Devonian and Mississippian rocks of central Arizona: U.S. Geological Survey Professional Paper 233-D, p. 67-112.
- Hunt, C.B., 1956, Cenozoic geology of the Colorado Plateau: U.S. Geological Survey Professional Paper 279, 99 p.
- Ijirigho, B.T., 1977, Pennsylvanian subsurface stratigraphy of the Black Mesa Basin and Four Corners area in northeastern Arizona: Tucson, Arizona, University of Arizona, unpublished M.S. thesis, 62 p.
- Jenney, J.P., and Stone, Claudia, eds., 1980, Studies in western Arizona: Arizona Geological Society Digest, V. XII, 322 p.
- Jenney, J.P., and Reynolds, S.J., eds., 1989, Geologic evolution of Arizona: Arizona Geological Society Digest 17, 866 p.
- Jordan, C.F., Jr., 1971, Lower Permian stratigraphy of southern New Mexico and West Texas: Houston, Texas, Rice University, unpublished Ph.D. dissertation, 136 p.
- Kashfi, M.S., 1983, Upper Devonian source and reservoir rocks in the Black Mesa Basin, northeastern Arizona: Oil and Gas Journal, v. 81, no. 50, December 12, p. 151-159.
- Kelley, V.C., and Clinton, N.J., 1960, Fracture systems and tectonic elements of the Colorado Plateau: University of New Mexico Publications in Geology no. 6, 104 p.
- Keith, S.B., and Wilt, J.C., 1986, Laramide orogeny in Arizona and adjacent regions--A strato-tectonic synthesis, *in* Beatty, B., and Wilkinson, P.A.K., eds., *Frontiers in geology and ore deposits of Arizona and the Southwest*: Arizona Geological Society Digest, V. XVI, p. 502-554.
- Kent, W.N., and Rawson, R.R., 1989, Depositional environments of the Mississippian Redwall Limestone in northeastern Arizona, *in* Fouch, T.D., and Magathan, E.R., eds., *Paleozoic paleogeography of west-central United States*: Society of Economic Paleontologists and Mineralogists, Symposium 1, p. 101-109.
- Kluth, C.F., 1986, Plate tectonics of the ancestral Rocky Mountains, *in* Peterson, J.A., ed., *Paleotectonics and sedimentation in the Rocky Mountain region, United States*: American Association of Petroleum Geologists Memoir 41, p. 353-369.

- Koons, E.D., 1945, Geology of the Uinkaret Plateau, northern Arizona: Geological Society of America Bulletin, v. 56, p. 151-180.
- Kottlowski, F.E., 1965, Sedimentary basins of south-central and south-western New Mexico: American Association of Petroleum Geologists Bulletin, v. 49, p. 2120-2139.
- Kottlowski, F.E., 1977, New Mexico geology and petroleum potential: Interstate Oil Compact Commission Committee Bulletin, v. 19, December, no. 2, p. 26-30.
- Kron, A., and Stix, J., 1982, Geothermal gradient map of the United States: National Oceanic and Atmospheric Administration, National Geophysics Data Center, prepared for Los Alamos National Laboratory, University of California, scale 1:2,500,000.
- Lessentine, R.H., 1965, Kaiparowits and Black Mesa Basin--Stratigraphic synthesis: American Association of Petroleum Geologists Bulletin, v. 49, p. 1997-2019.
- Lochman-Balk, C., 1972, Cambrian System, *in* Mallory, W.W., ed., Geologic atlas of the Rocky Mountain region: Rocky Mountain Association of Geologists, p. 60-75.
- Lovejoy, E.M.P., 1973, Major Early Cenozoic deformation along Hurricane Fault zone, Utah and Arizona: American Association of Petroleum Geologists Bulletin, v. 57, p. 510-519.
- Luedke, R.G., and Smith, R.L., 1978, Map showing distribution, composition, and age of Late Cenozoic volcanic centers in Arizona and New Mexico: U.S. Geological Survey Miscellaneous Investigations Series Map I-1091-A, scale 1:1,000,000.
- Lyth, A.L., Jr., 1971, Summary of possible future petroleum potential, region 3, Western Rocky Mountains, *in* Cram, I.H., ed., Future petroleum provinces of the United States--Their geology and potential: American Association of Petroleum Geologists Memoir 15, vol. 1, p. 406-412.
- Mack, G.H., Lawton, T.F., and Lucas, S.G., eds., 1988, Cretaceous and Laramide tectonic evolution of southwestern New Mexico: New Mexico Geological Society, 39th Field Conference Guidebook, 216 p.
- McKee, E.D., 1967, Arizona and western New Mexico, *in* Paleotectonic investigations of the Permian System in the United States: U.S. Geological Survey Professional Paper 515-J, p. 200-223.
- McKee, E.D., 1975, Arizona, *in* McKee, E.D., and Crosby, E.J., eds., Paleotectonic investigations of the Pennsylvanian System in the United States: U.S. Geological Survey Professional Paper 853-P, p. 293-309.

- McKee, E.D., 1979, Arizona, *in* Craig, L.C., and Conner, C.W., coordinators, Paleotectonic investigations of the Mississippian System in the United States, pt. 1: U.S. Geological Survey Professional Paper 1010-L, p. 198-207.
- McKee, E.D., 1982, The Supai Group of Grand Canyon: U.S. Geological Survey Professional Paper 1173, 504 p.
- McKenny, J.W., and Masters, J.A., 1968, Dineh-bi-Keyah Field, Apache County, Arizona: American Association of Petroleum Geologists Bulletin, v. 52, p. 2045-2057.
- Mallory, W.W., ed., 1972, Geologic atlas of the Rocky Mountain region: Rocky Mountain Association of Geologists, 331 p.
- Muehlberger, W.R., 1980, Texas lineament revisited, *in* Dickerson, P.W., and Hoffer, J.M., eds., Trans-Pecos region guidebook: New Mexico Geological Society 31st Field Conference, p. 113-121.
- Nations, J.D., Brennan, D.J., and Ybarra, R.A., 1989, Oil and gas in Arizona, *in* Jenney, J.P., and Reynolds, S.J., eds., Geologic evolution of Arizona: Arizona Geological Society Digest, V. 17, p. 795-816.
- Nations, J.D., Conway, C.M., and Swann, G.A., eds., 1986, Geology of central and northern Arizona: Geological Society of America Field Trip Guidebook for Rocky Mountain Section, Flagstaff, Arizona, Northern Arizona University, Geology Department, 176 p.
- Nations, J.D., Doss, A.K., and Ybarra, R.A., 1983a, Geologic setting of oil and gas exploration in Arizona, *in* Fassett, J.E., ed., Oil and gas fields of the Four Corners area, V. III: Four Corners Geological Society, p. 763-772.
- Nations, J.D., Doss, A.K., and Ybarra, R.A., 1983b, Stratigraphy and oil and gas production of Arizona, 1978-1983, *in* Fassett, J.E., ed., Oil and gas fields of the Four Corners area, V. III: Four Corners Geological Society, p. 832-843.
- Nations, J.D., and Eaton, J.G., eds., 1991, Stratigraphy, depositional environments, and sedimentary tectonics of the western margin, Cretaceous western Interior Seaway: Geological Society of America Special Paper 260.
- Nations, J.D., Landye, J.J., and Hevly, R.H., 1982, Location and chronology of Tertiary sedimentary deposits in Arizona--A review, *in* Ingersoll, R.V., and Woodburne, M.O., eds., Cenozoic nonmarine deposits of California and Arizona: Society of Economic Paleontologists and Mineralogists, Pacific Coast Section Symposium, p. 107-122.

- Nations, J.D., and Ranney, W.D.R., 1989, Geologic model of a Cenozoic basin in Arizona, *in* Tooker, E.W., ed., Arizona's industrial rock and mineral resources--Workshop proceedings: U.S. Geological Survey Bulletin 1905, [in cooperation with Arizona Geological Survey], p. 44-53.
- Nations, J.D., and Stump, E., 1981, Geology of Arizona: Dubuque, Iowa, Kendall Hunt Publishing Company, 210 p.
- Nations, J.D., Wilt, J.C., and Hevly, R.H., 1985, Cenozoic paleogeography of Arizona, *in* Flores, R.M., and Kaplan, S.S., eds., Cenozoic paleogeography of the west-central United States: Society of Economic Paleontologists and Mineralogists, Rocky Mountain Paleogeography Symposium 3, p. 335-355.
- Oil and Gas Journal, 1979, Arizona area offers good potential for new finds: Oil and Gas Journal, v. 77, no. 44, p. 203-204.
- Olmsted, R.H., Loeltz, O.J., and Irelan, B., 1973, Geohydrology of the Yuma area, Arizona and California: U.S. Geological Survey Professional Paper 486-H, 227 p.
- Oppenheimer, J.M., and Sumner, J.S., 1980, Depth-to-bedrock map, Basin and Range Province, Arizona: Tucson, Arizona, University of Arizona, Department of Geosciences, scale 1:1,000,000.
- O'Sullivan, R.B., 1969, Crude oil, natural gas, and other associated gases, *in* Mineral and water resources of Arizona: Arizona Bureau of Mines Bulletin 180, p. 70-91.
- Peirce, H.W., 1976a, Tectonic significance of Basin and Range thick evaporite deposits: Arizona Geological Society Digest V. X, p. 325-339.
- Peirce, H.W., 1976b, Elements of Paleozoic tectonism in Arizona: Arizona Geological Society Digest V. X, p. 37-57.
- Peirce, H.W., 1979, The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States--Arizona: U.S. Geological Survey Professional Paper 1110-Z, 20 p.
- Peirce, H.W., 1981, Major Arizona salt deposits: Arizona Bureau of Geology and Mineral Technology Fieldnotes, v. 11, no. 4, p. 1-4.
- Peirce, H.W., 1982, The search for petroleum in Arizona: Arizona Bureau of Geology and Mineral Technology, Fieldnotes, v. 12, no. 2, p. 1-5.
- Peirce, H.W., and Gerrard, T.A., 1966, Evaporites of the Permian Holbrook Basin, Arizona, *in* Rau, J.L., ed., Second symposium on salt: Northern Ohio Geological Society, v. 1, p. 1-10.

- Peirce, H.W., Jones, Nile, and Rogers, Ralph, 1977, A survey of uranium favorability of Paleozoic rocks in the Mogollon Rim and Slope region east-central Arizona: Arizona Bureau of Geology and Mineral Technology Circular 19, 70 p.
- Peirce, H.W., Keith, S.B., and Wilt, J.C., 1970 (reprinted 1975), Coal, oil, natural gas, helium, and uranium in Arizona: Arizona Bureau of Mines Bulletin 182, 289 p. and 19 plates.
- Peirce, H.W., and Scurlock, J.R., 1972, Arizona well information: Arizona Bureau of Mines Bulletin 185, 195 p.
- Peterson, J.A., ed., 1986, Paleotectonics and sedimentation in the Rocky Mountain region, United States: American Association of Petroleum Geologists Memoir 41, 693 p.
- Plauchut, Bernard, Tardy, Marc, Blanchet, Rene, Zimmermann, Matthis, Cabezas, Pascal, Rosaz, Thierry, and Sosson, Marc, 1989, Etude Geologique des Cordilleres Nord-Americaines; Les lineaments du Texas et Caltam entre Cordilleres Americanines et Sierras Madres Mexicaines--Nature, origine et evolution structurale; Les montagnes rocheuses meridionales au centre-ouest du Nouveau-Mexique; Le passage des Cordilleres Nord-Americaines aux Sierras Madres Mexicaines le long du Texas lineament geologie du SW du Nouveau Mexique; Tectoniques superposees dans le sud-est de l'Arizona: New Mexico Bureau of Mines and Mineral Resources Open-File Report 385, 76 p.
- Poole, F.G., and 8 others, 1967, Devonian of the Southwestern United States, *in* Oswald, D.H., ed., International Symposium on the Devonian System: Society of Petroleum Geologists, Proceedings, v. 1, September 1967, Calgary, Canada, p. 879-912.
- Pray, L.C., 1977, Stratigraphic and structural features of the Sacramento Mountain escarpment, New Mexico, *in* Butler, J.H., ed., Geology of the Sacramento Mountains, Otero County, New Mexico: West Texas Geological Society Guidebook, Publication no. 1977-68, p. 73-89.
- Pye, W.D., 1967, Arizona--A new frontier to explore: Oil and Gas Journal, v. 65, May 8th, p. 168-172.
- Pyron, A.J., and Gray, J.H., 1985, Don't ignore the Orogrande; its potential is promising: World Oil, May, p. 59-67.
- Rauzi, S.L., 1990, Distribution of Proterozoic hydrocarbon source rock in northern Arizona and southern Utah: Phoenix, Arizona, Arizona Oil and Gas Conservation Commission Special Publication 5, 38 p.; map scale 1:500,000.

- Rauzi, S.L., 1992, Exploration well tests salt near Phoenix [Arizona], *in* Arizona Geology: Arizona Geological Survey, v. 22, no. 4, p. 910.
- Rauzi, S.L., 1993, Oil and natural gas occurrence in Arizona: Arizona Geological Survey Publication OG-35 (OGCC Chart C-1).
- Rauzi, S.L., 1994, Implications of live oil shows in eastern Arizona geothermal test: Arizona Geological Survey Open-File Report 94-1, 16 p; 12 maps, scale 1:500,000.
- Reif, D.M., and Robinson, J.P., 1981, Geophysical, geochemical, and petrographic data and regional correlation from the Arizona State A-1 well, Pinal County, Arizona: Arizona Geological Society Digest, V. XIII, p. 99-109.
- Reiter, M., Mansure, A.J., and Shearer, C., 1979, Geothermal characteristics of the Colorado Plateau: Tectonophysics, v. 61, p. 183-195.
- Reynolds, M.W., Palacas, J.G., and Elston, D.P., 1988, Potential petroleum source rocks in the Late Proterozoic Chuar Group, Grand Canyon, Arizona, *in* Carter, L.M.H., ed., USGS research on energy resources--1988, Program and abstracts: U.S. Geological Survey Circular 1025, p. 49-50.
- Reynolds, S.J., 1988, Geologic map of Arizona: Arizona Geological Survey Map 26, scale 1:1,000,000.
- Reynolds, S.J., Roddy, M.S., and Welty, J.W., 1986, Map of Paleozoic rocks and conodont alteration indices in Arizona: Arizona Bureau of Geology and Mineral Technology Open-File Report 86-1, scale 1:1,000,000.
- Reynolds, S.J., and Trapp, R.A., 1986, Computerized databases and database access programs of the Arizona Geological Survey: Arizona Geological Survey Open-File Report 86-7, 69 p.
- Reynolds, S.J., Welty, J.W., and Spencer, J.E., 1986, Volcanic history of Arizona: Arizona Bureau of Geology and Mineral Technology Field Notes, v. 16, no. 2, p. 1-5.
- Rieker, R.E., ed., 1979, Rio Grande Rift--Tectonics and magmatism: Washington, D.C., American Geophysical Union, 438 p.
- Robertson, J.M., and Broadhead, R.F., project directors, 1993, Atlas of major Rocky Mountain gas reservoirs: New Mexico Bureau of Mines and Mineral Resources (published for a consortium of agencies including Gas Research Institute, U.S. Department of Energy, Colorado Geological Survey, Utah Geological Survey, and Geological Survey of Wyoming), 206 p. (oversize).



- Ross, C.A., and Ross, J.R.P., 1986, Paleozoic paleotectonics and sedimentation in Arizona and New Mexico, *in* Peterson, J.A., ed., Paleotectonics and sedimentation in the Rocky Mountain region, United States: American Association of Petroleum Geologists Memoir 41, p. 653-668.
- Roybal, G.H., 1981, Facies development in a Lower Cretaceous coral-rudist patch reef (Mural Limestone, southeast Arizona): Rocky Mountain Association of Geologists, *The Mountain Geologist*, v. 18, no. 3, p. 46-56.
- Ryder, R.T., 1983a, Petroleum potential of wilderness lands in Arizona, *in* Miller, B.M., ed., Petroleum potential of wilderness lands in the western United States: U.S. Geological Survey Circular 902-C, p. 1-22.
- Ryder, R.T., 1983b, Petroleum potential of wilderness lands in New Mexico, *in* Miller, B.M., ed., Petroleum potential of wilderness lands in the Western United States: U.S. Geological Survey Circular 902-I, p. 1-33.
- Schmidt, Nancy, Reynolds, S.J., and Horstman, K.C., 1991, AZGEOBIB--A preliminary list of references on the geology of Arizona: Arizona Geological Survey Open-File Report 91-4, 302 p.
- Scurlock, J.R., 1971, Geologic structure map of northern Arizona: Phoenix, Arizona, Arizona Oil and Gas Conservation Commission, scale 1:375,000.
- Seager, W.R., 1983, Laramide wrench faults, basement-cored uplifts, and complementary basins in southern New Mexico: New Mexico Bureau of Mines and Mineral Resources, *New Mexico Geology*, v. 5, no. 4, p. 69-76.
- Seager, W.R., Clemons, R.E., and Callender, J.F., eds., 1975, Guidebook of the Las Cruces country: New Mexico Geological Society, 26th Field Conference, 345 p.
- Seager, W.R., and Morgan, P., 1979, Rio Grande Rift in southern New Mexico, West Texas, and northern Chihuahua, *in* Riecker, R.E., ed., Rio Grande Rift--Tectonism and magmatism: Washington, D.C., American Geophysical Union, p. 87-106.
- Shride, A.F., 1967, Younger Precambrian geology in southern Arizona: U.S. Geological Survey Professional Paper 566, 89 p.
- Smith, R.B., and Eaton, G.P., eds., 1978, Cenozoic tectonics and regional geophysics of the western Cordillera: Geological Society of America Memoir 152, 388 p.
- Stewart, W.J., ed., 1969, Delaware Basin exploration guidebook: West Texas Geological Society Publication no. 68-55a, 170 p.

- Stone, Claudia, and Jenney, J.P., eds., 1981: Arizona Geological Society Digest, V. XIII, 193 p.
- Stone, Claudia, and Witcher, J.C., 1982, Geothermal energy in Arizona--Final contract report: Arizona Bureau of Geology and Mineral Technology for the U.S. Department of Energy, Contract No. DE-FC07-79ID12009, 398 p.
- Sumner, J.S., 1975, Bouguer gravity map of Arizona: Tucson, Arizona, University of Arizona Laboratory of Geophysics, scale 1:1,000,000.
- Sumner, J.S., 1985, Crustal geology of Arizona as interpreted from magnetic, gravity, and geologic data, *in* Hinze, W.J., ed., The utility of regional gravity and magnetic anomaly maps: Society of Exploration Geophysicists, p. 164-180.
- Sylvester, A.G., comp., 1984, Wrench fault tectonics: American Association of Petroleum Geologists Reprint Series no. 23, 374 p.
- Teichert, C., 1965, Devonian rocks and paleogeography of central Arizona: U.S. Geological Survey Professional Paper 464, 181 p.
- Terra Graphics, 1977, Oil and gas production map of the western United States: Denver, Colorado, Terra Graphics, scale approximately 1:3,250,000.
- Thompson, G.A., and Zoback, M.L., 1979, Regional geophysics of the Colorado Plateau: Tectonophysics, v. 61, p. 149-181.
- Thompson, Sam, III, 1981, Analyses of petroleum source and reservoir rocks in southwestern New Mexico--Petroleum source rocks in exploration wells drilled to Paleozoic or Mesozoic units, Hidalgo and Grant Counties, New Mexico: New Mexico Bureau of Mines and Mineral Resources Open-File Report OF-153, 120 p.
- Thompson, Sam, III, 1982, Oil and gas exploration wells in southwestern New Mexico, *in* Powers, R.B., ed., Geologic studies of the Cordilleran thrust belt, v. II: Rocky Mountain Association of Geologists, p. 521-535.
- Thompson, Sam, III, and Bierberman, R.A., 1975, Oil and gas exploration wells in Doña Ana County, New Mexico, *in* Seager, W.R., Clemons, R.E., and Callender, J.F., eds., Guidebook of the Las Cruces Country: New Mexico Geological Society, 26th Field Conference, p. 171-174.
- Thompson, Sam, III, and Jacka, A.D., 1981, Pennsylvanian stratigraphy, petrography, and petroleum geology of Big Hatchet Peak section, Hidalgo County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Circular 176, 129 p.

- Titley, S.R., ed., 1968, Southern Arizona Guidebook III: Arizona Geological Society (in cooperation with the Geological Society of America), 354 p.
- Tooker, E.W., comp.-ed., 1989, Arizona's industrial rock and mineral resources-- Workshop proceedings: U.S. Geological Survey Bulletin 1905 (with the Arizona Geological Survey), 93 p.
- Tovar R, J.C., 1981, Provincias con posibilidades petroleras en el Distrito de Chihuahua: Boletín de la Asociación Mexicana de Geólogos Petroleros, v. 33, no. 1, p. 25-51.
- Tucker, M.L., 1983, Geochemical correlation of crude oil and source rocks in Black Mesa Basin, northeastern Arizona: Flagstaff, Arizona, Northern Arizona University, unpublished M.S. thesis, 80 p.
- Turner, D.S., 1968, Natural gas in Black Mesa Basin, northeastern Arizona, *in* Beebe, W., ed., Natural gases of North America: American Association of Petroleum Geologists Memoir 9, v. 2, p. 1357-1370.
- Wardlaw, B.R., and Harris, A.G., 1984, Conodont-based thermal maturation of Paleozoic rocks in Arizona: American Association of Petroleum Geologists Bulletin, v. 69, p. 1101-1106.
- Weiss, Emanuel, 1991, Regional ground-water flow in upper and middle Paleozoic rocks in southeastern Utah and adjacent parts of Arizona, Colorado, and New Mexico: U.S. Geological Survey Water-Resources Investigations Report 90-4079, 57 p.
- Wengerd, S.A., 1970, Petroleum prospects in southwesternmost New Mexico, *in* Woodward, L.A., ed., Guidebook of the Tyrone--Big Hatchet Mountains Florida Mountains region: New Mexico Geological Society, 21st Field Conference, p. 91-104.
- Wilson, E.D., 1962, A r̄sum̄ of the geology of Arizona: Arizona Bureau of Mines Bulletin 171, 140 p.
- Wilson, E.D., and Moore, R.T., 1962, Geologic cross-sections of Arizona 1-8: Arizona Bureau of Mines, horizontal scale 1 inch equals 3 miles, vertical scale 1 inch equals 500 feet.
- Wilson, E.D., Moore, R.T., and Cooper, J.R., 1969, Geologic map of Arizona: Arizona Bureau of Mines and U.S. Geological Survey, scale 1:500,000.
- Wilson, J.L., 1989, Lower and Middle Pennsylvanian strata in the Orogrande and Pedregosa Basins, New Mexico: New Mexico Bureau of Geology and Mineral Resources Bulletin 124, 16 p.

- Wilt, J.C., and Jenney, J.P., eds., 1976, Tectonic digest: Arizona Geological Society digest, V. X, 430 p.; 4 plates.
- Woodward-Clyde Consultants, 1983, Overview of the regional geology of the Paradox Basin study region, *for* the U.S. Department of Energy, Office of Nuclear Waste Isolation, ONWI-92, distribution category UC-70: San Francisco, California, 433 p.
- Woodward, L.A., ed., 1970, Guidebook of the Tyrone--Big Hatchet Mountains Florida Mountains region: New Mexico Geological Society, 21st Field Conference, 162 p.
- Woodward, L.A., 1973, Structural framework and tectonic evolution of the Four Corners region of the Colorado Plateau, *in* James, H.L., ed., Guidebook of Monument Valley and vicinity Arizona and Utah: New Mexico Geological Society, 24th Field Conference, p. 94-98.
- Woodward, L.A., comp., 1984, Tectonic map of the Rocky Mountain region of the United States, *in* Sloss, L.L., ed., Sedimentary cover--North American craton: Geological Society of America Decade of North American Geology Project, v. D-2, plate 2, scale 1:2,500,000.
- Woodward, L.A., and Duchene, H.R., 1981, Overthrust belt of southwestern New Mexico: Comparison with Wyoming-Utah overthrust belt: American Association of Petroleum Geologists Bulletin, v. 65, p. 722-729.
- Woodward, L.A., and Grant, P.R., 1986, Central-western New Mexico--An exploration frontier for oil and gas, *in* Clemons, R.E., King, W.E., and Mack, G.H., eds., Truth or Consequences region: New Mexico Geological Society Guidebook, 37th Field Conference, p. 307-314.
- Woodward, L.A., and Northrop, S.A., eds., 1976, Tectonics and mineral resources of Southwestern North America: New Mexico Geological Society Special Publication no. 6, 218 p.
- Wrucke, C.T., and Armstrong, A.K., 1987, Paleozoic stratigraphic section in Dry Canyon, Whetstone Mountains, Cochise County, Arizona, *in* Hill, M.L., ed., Centennial field guide volume 1: Geological Society of America, Cordilleran Section, p. 29-34.

- Wrucke, C.T., Otton, J.K., and Desborough, G.A., 1986, Summary and origin of the mineral commodities in the Middle Proterozoic Apache Group in central Arizona, *in* Geatty, B., and Wilkinson, P.A.K., eds., *Frontiers in geology and ore deposits of Arizona and the Southwest: Arizona Geological Society Digest*, V. XVI, p. 12-17.
- Young, R.A., 1979, Laramide deformation, erosion, and plutonism along the southwestern margin of the Colorado Plateau: *Tectonophysics*, v. 61, p. 25-47.
- Zietz, I., comp., 1982, Composite magnetic anomaly map of the United States: U.S. Geological Survey and Society of Exploration Geophysicists, scale 1:2,500,000.